

# Developing a collaborative approach to monitor drug related deaths helps to improve care for people in treatment


“The benefits of a regional collaborative drug related death surveillance system: Insights from development of a monitoring system in the North West of England” Mark Whitfield, Howard Reed

**Drug Related Deaths (DRDs)** across England and Wales are at their highest level since records began in 1993, with deaths from overdose being particularly prominent in recent years. Across Cheshire and Merseyside, a region in the north west of England, deaths are also at their highest level since records began. There may be many reasons for this including increased purity levels of street drugs, the increasing age of people accessing Treatment, and service cuts due to austerity.

## Drug fatalities highest where treatment cutbacks deepest

Reducing funds for councils to spend on addiction services 'catalyst for disaster' warn academics

The Guardian, 14<sup>th</sup> October 2017




**In response**, seven of the nine local authorities within Cheshire and Merseyside region established a review process to examine DRDs in order to explore common themes, identify recurring issues and to share findings. The monitoring system was developed by Liverpool John Moores University's Public Health Institute (PHI) who collate information from various sources and who chair regular panels. The process is based on recommendations from the Local Government Association<sup>(1)</sup> and Public Health England<sup>(2)</sup>.

<sup>(1)</sup> "Preventing drug related deaths" Local Government Association, 2017, <sup>(2)</sup> "Understanding and Preventing Drug Related Deaths", PHE, 2016



## HOW DOES IT WORK



### WHO ATTENDS THE PANELS?

- Treatment service staff
- Consultant prescriber
- Local authority public health commissioner
- Social services staff
- Hostels
- COPD (Respiratory Care)
- End of Life Care
- Medicines Management




### Common Themes

- **Individuals** tend to live alone which Naloxone may not be useful for
- **A high volume** of medication is prescribed by GP's sometimes in excess of 20 per individual
- **Increasing numbers** of older people accessing treatment have high levels of COPD/respiratory disease
- **People not in treatment** are dying at a younger age
- **Sup-optimal prescribing** of methadone is common, but there may be clinical reasons for this
- **Depression** is the highest commonly identified mental health condition

### Some actions resulting from DRD panels

- ✓ Pathways have been developed between drug treatment services and other services including palliative care and prison services
- ✓ Local protocols reviewed around bridging prescriptions
- ✓ Day programme in signing skills for staff was developed
- ✓ Awareness package for staff in COPD symptoms to give them greater confidence
- ✓ Focus on individuals living alone with risk assessments carried out
- ✓ Exploration of offer for individuals not engaging with treatment
- ✓ Liaison with nursing homes and sheltered accommodation for older people
- ✓ Campaign around smoking cessation implemented for treatment service staff
- ✓ Links strengthened with services assisting both families and services users through bereavement
- ✓ Recovery position training delivered to staff

**Implementation barriers** included data sharing issues (particularly with coroner data), funding for the system during a time of significant pressures on public health budgets, getting the key stakeholders together at the same time and dealing with the volume of deaths within the available time



Take a picture to download this poster

[ims.ljmu.ac.uk/posters](https://ims.ljmu.ac.uk/posters)